

Chen Inequalities for Submanifolds of Real Space Forms with a Ricci Quarter-Symmetric Metric Connection

Nergiz (Önen) Poyraz* and Halil İbrahim Yoldaş

(Communicated by H. Hilmi Hacısalihoğlu)

ABSTRACT

In this paper, we establish some inequalities for submanifolds of real space forms endowed with a Ricci quarter-symmetric metric connection. Using these inequalities, we obtain the relation between Ricci curvature, scalar curvature and the mean curvature endowed with the Ricci quarter-symmetric metric connection.

Keywords: Chen inequality; Ricci quarter-symmetric metric connection; Ricci curvature.

AMS Subject Classification (2010): 53B05 ; 53B15; 53C40.

1. Introduction

In [8], the idea of a Ricci quarter-symmetric metric connection on a Riemannian manifold was introduced and presented by Kamilya and De. They also found necessary and sufficient conditions for the symmetry of the Ricci tensor of a Ricci quarter-symmetric metric connection and showed that conformal curvature tensor of induced connection ∇ and linear connection $\tilde{\nabla}$ are equal [8]. Before this work, a few papers had been written about the studies of various types of a quarter-symmetric metric connection and their properties in [11] and [12].

In 1993, Chen [4] introduced a new Riemannian invariant for a Riemannian manifold M as follows:

$$\delta_M = \tau(p) - \inf(K)(p), \quad (1.1)$$

where $\tau(p)$ is scalar curvature of M and

$$\inf(K)(p) = \inf\{K(\Pi) : K(\Pi) \text{ is a plane section of } T_p M\}.$$

Chen gave the following general optimal inequality involving the new intrinsic invariant δ_M , the squared mean curvature $\|H\|^2$ for an n -dimensional submanifold M in a real space form $R(c)$ of constant sectional curvature c :

$$\delta_M \leq \frac{n^2(n-2)}{2(n-2)} \|H\|^2 + \frac{1}{2}(n+1)(n-2)c. \quad (1.2)$$

[3].

Also, Chen established a sharp inequality between the main intrinsic curvatures (the sectional curvature and the scalar curvature) and the main extrinsic curvatures (the squared mean curvature) for a submanifold in real space form $R^m(\bar{c})$, well-known as *Chen inequalities*, in [2] as follows:

For each unit tangent vector $X \in T_p M^n$,

$$H^2(p) \geq \frac{4}{n^2} \{Ric(X) - (n-1)\bar{c}\}, \quad (1.3)$$

where H^2 is the squared mean curvature and $Ric(X)$ is Ricci curvature of M^n at X .

In [7], Hong and Tripathi presented a general inequality for submanifolds of a Riemannian manifold by using (1.3). In [13], this inequality was named Chen-Ricci inequality by Tripathi. In fact, the general inequality obtained in [7] is a special case of Theorem 3.1 of [5]. Later, Mihai and Özgür in [10] proved Chen inequalities for submanifolds of real space forms endowed with a semi-symmetric metric connection. Moreover, several works in this direction is studied [1, 6, 9, 13].

The paper is organized as follows: Section 1 is concerned with introduction. In section 2, we give some basic concepts on submanifolds of Riemannian manifold endowed with Ricci quarter-symmetric metric connection which will be used throughout this paper. In section 3, we find some inequalities for submanifolds of real space forms endowed with a Ricci quarter-symmetric metric connection. Considering these inequalities, we obtain the relation between Ricci curvature, scalar curvature and the mean curvature endowed with the Ricci quarter-symmetric metric connection.

2. Preliminaries

Let \widetilde{M} be an m -dimensional Riemannian manifold and $\widetilde{\nabla}$ a linear connection on \widetilde{M} . A linear connection $\widetilde{\nabla}$ is said to be Ricci quarter-symmetric connection if the torsion tensor \widetilde{T} is of the form

$$\widetilde{T}(X, Y) = \pi(Y)LX - \pi(X)LY, \tag{2.1}$$

where $\widetilde{\pi}$ is a 1-form and L is the (1, 1) Ricci tensor defined by

$$\widetilde{g}(LX, Y) = S(X, Y) \tag{2.2}$$

S is the Ricci tensor of \widetilde{M} .

A linear connection $\widetilde{\nabla}$ is called a metric connection if

$$(\widetilde{\nabla}_X \widetilde{g})(Y, Z) = 0. \tag{2.3}$$

Following [8], a Ricci quarter-symmetric metric connection $\widetilde{\nabla}$ on \widetilde{M} is given by

$$\widetilde{\nabla}_{\widetilde{X}} \widetilde{Y} = \overset{\circ}{\nabla}_{\widetilde{X}} \widetilde{Y} + \pi(\widetilde{Y})L\widetilde{X} - S(\widetilde{X}, \widetilde{Y})P \tag{2.4}$$

for any vector fields \widetilde{X} and \widetilde{Y} of \widetilde{M} , where $\overset{\circ}{\nabla}$ denotes the Levi-Civita connection with respect to the Riemannian metric \widetilde{g} , π is a 1-form and P is the vector field defined by

$$\widetilde{g}(P, \widetilde{X}) = \pi(\widetilde{X})$$

for an arbitrary vector field \widetilde{X} of \widetilde{M} .

From now on, we will consider a Riemannian manifold \widetilde{M} endowed with a Ricci quarter-symmetric metric connection $\widetilde{\nabla}$ and the Levi-Civita connection denoted by $\overset{\circ}{\nabla}$.

Let M^n be an n -dimensional submanifold of an m -dimensional Riemannian manifold \widetilde{M} . On the submanifold M^n we consider the induced Ricci quarter-symmetric metric connection denoted by ∇ and the induced Levi-Civita connection denoted by $\overset{\circ}{\nabla}$.

Let \widetilde{R} be the curvature tensor of \widetilde{M} with respect to $\widetilde{\nabla}$ and $\overset{\circ}{R}$ the curvature tensor of \widetilde{M} with respect to $\overset{\circ}{\nabla}$. We also denote by R and $\overset{\circ}{R}$ the curvature tensors of ∇ and $\overset{\circ}{\nabla}$, respectively, on M .

The Gauss formulas with respect to ∇ and $\overset{\circ}{\nabla}$, respectively, can be written as:

$$\widetilde{\nabla}_X Y = \nabla_X Y + h(X, Y), \tag{2.5}$$

$$\overset{\circ}{\nabla}_X Y = \overset{\circ}{\nabla}_X Y + \overset{\circ}{h}(X, Y), \tag{2.6}$$

where $\overset{\circ}{h}$ is the second fundamental form of M in \widetilde{M} and h is a (0, 2)-tensor on M .

For any orthonormal basis $\{e_1, \dots, e_n\}$ of the tangent space T_pM^n , the mean curvature vector $H(p)$ is given by

$$H(p) = \frac{1}{n} \sum_{i=1}^n h(e_i, e_i).$$

If $h = 0$ (respectively $H = 0$), then the submanifold M^n is called totally geodesic (minimal) in \widetilde{M} . If $h(X, Y) = g(X, Y)H$ for all $X, Y \in TM$, then M^n is said to be totally umbilical.

The Gauss equation with respect to the Ricci quarter-symmetric metric connection is

$$R(X, Y, Z, W) = \widetilde{R}(X, Y, Z, W) + g(h(X, Z), h(Y, W)) - g(h(X, W), h(Y, Z)). \quad (2.7)$$

The curvature tensor $\overset{\circ}{R}$ with respect to the Levi-Civita connection $\overset{\circ}{\nabla}$ on \widetilde{M} (e) is expressed by

$$\overset{\circ}{R}(X, Y, Z, W) = c\{g(X, W)g(Y, Z) - g(Y, W)g(X, Z)\}. \quad (2.8)$$

Let $\Pi = \text{Span}\{e_i, e_j\}$ be 2-dimensional non-degenerate plane of the tangent space T_pM at $p \in M$. Then the number

$$K_{ij} = \frac{g(R(e_j, e_i)e_i, e_j)}{g(e_i, e_i)g(e_j, e_j) - g(e_i, e_j)^2} \quad (2.9)$$

is called the sectional curvature of the section Π at $p \in M$.

Let M^n be an n -dimensional Riemannian manifold. We denote by $K(\pi)$ the sectional curvature of M^n associated with a plane section $\pi \subset T_pM^n$, $p \in M^n$. If $\{e_1, \dots, e_n\}$ is an orthonormal basis of the tangent space T_pM^n , then the scalar curvature τ at p is defined by

$$\tau(p) = \sum_{1 \leq i < j \leq n} K_{ij}.$$

Let M^n be an n -dimensional Riemannian manifold, L be a k -plane section of T_pM^n , $p \in M^n$, and X be a unit vector in L . We choose an orthonormal basis $\{e_1, \dots, e_k\}$ of L such that $e_1 = X$.

One defines [2] the Ricci curvature (or k -Ricci curvature) of L at X by

$$Ric_L(X) = K_{12} + K_{13} + \dots + K_{1k},$$

where K_{ij} denotes, as usual, the sectional curvature of the 2-plane section spanned by e_i, e_j . For each integer k , $2 \leq k \leq n$, the Riemannian invariant θ_k on M^n is defined by:

$$\theta_k(p) = \frac{1}{k-1} \inf_{L, X} Ric_L(X), \quad p \in M, \quad (2.10)$$

where L runs over all k -plane sections in T_pM^n and X runs over all unit vectors in L .

Then the curvature tensor \widetilde{R} with respect to the Ricci quarter-symmetric metric connection $\widetilde{\nabla}$ on \widetilde{M} can be shown that [8]

$$\begin{aligned} \widetilde{R}(X, Y)Z &= \overset{\circ}{R}(X, Y)Z - M(Y, Z)LX + M(X, Z)LY \\ &\quad - S(Y, Z)QX + S(X, Z)QY + \pi(Z)[(\overset{\circ}{\nabla}_X L)Y - (\overset{\circ}{\nabla}_Y L)X] \\ &\quad - [(\overset{\circ}{\nabla}_X S)(Y, Z) - (\overset{\circ}{\nabla}_Y S)(X, Z)]P, \end{aligned} \quad (2.11)$$

where M is tensor field of type $(0, 2)$ defined by

$$M(X, Y) = g(QX, Y) = (\overset{\circ}{\nabla}_X \pi)Y - \pi(Y)\pi(LX) + \frac{1}{2}\pi(P)S(X, Y) \quad (2.12)$$

and Q is a tensor field of type $(2, 1)$ defined by

$$QX = \overset{\circ}{\nabla}_X P - \pi(LX)P + \frac{1}{2}\pi(P)LX. \quad (2.13)$$

Here we shall consider M^n to be an Einstein manifold, that is,

$$S(X, Y) = \frac{\overset{\circ}{\tau}}{n}g(X, Y), \tag{2.14}$$

where $\overset{\circ}{\tau}$ is the scalar curvature.

Throughout this paper, we assume that M^n is an Einstein manifold.

Considering (2.11) and (2.14) we get

$$\tilde{R}(X, Y)Z = \overset{\circ}{R}(X, Y)Z - \frac{\overset{\circ}{\tau}}{n}[M(Y, Z)X - M(X, Z)Y + g(Y, Z)QX - g(X, Z)QY]. \tag{2.15}$$

Contracting (2.15) with respect to X , we get

$$\tilde{S}(Y, Z) = \frac{\overset{\circ}{\tau}}{n}[g(Y, Z) - \{(n - 2)M(Y, Z) + mg(Y, Z)\}], \tag{2.16}$$

where \tilde{S} is the Ricci tensor of $\tilde{\nabla}$ and m is the trace of $M(Y, Z)$. Now putting $Y = Z = e_i$, where $\{e_1, \dots, e_n\}$ is an orthonormal basis of the tangent space at any point, we get by taking the sum for $1 \leq i \leq n$ in the relation (2.16)

$$\tilde{\tau} = \frac{\overset{\circ}{\tau}}{n}[n - 2(n - 1)m], \tag{2.17}$$

where $\tilde{\tau}$ is the scalar curvature of $\tilde{\nabla}$.

3. k-Ricci Curvature and k-Scalar Curvature

In this section, a sharp relation between the Ricci curvature in the direction of unit tangent vector X and the mean curvature H with respect to Ricci quarter-symmetric metric connection $\tilde{\nabla}$ is established. Using this inequality, a relationship between the k -Ricci curvature of M^n and the squared mean curvature $\|H\|^2$ is showed. From now on, we assume that the vector field P is tangent to M^n .

Denote by

$$N(p) = \{X \in T_pM^n \mid h(X, Y) = 0, \forall Y \in T_pM^n\}.$$

Theorem 3.1. *Let M^n be an n -dimensional submanifold of an m -dimensional real space form $\tilde{M}(c)$ of constant sectional curvature c endowed with Ricci quarter-symmetric metric connection $\tilde{\nabla}$. Then, the following statements are true.*

(i) *For each unit vector $X \in T_pM^n$ we have*

$$Ric(X) \leq \frac{1}{4}n^2 \|H\|^2 - (n - 1)c[1 - m + (n - 2)M(X, X)], \tag{3.1}$$

where m is the trace of M .

(ii) *The equality case of (3.1) is satisfied by unit vector $X \in T_pM^n$ if and only if*

$$\begin{aligned} h(X, Y) &= 0, \text{ for all } Y \in T_pM^n \text{ orthogonal to } X, \\ h(X, X) &= \frac{n}{2}H(p). \end{aligned} \tag{3.2}$$

(iii) *The equality case of (3.1) holds for all unit vector $X \in T_pM^n$ if and only if either p is a totally geodesic point or $n = 2$ and p is a totally umbilical point.*

Proof. From (2.7) and (2.15) we get

$$2\tau(p) = n(n - 1)c - 2(n - 1)^2cm + n^2 \|H\|^2 - \|h\|^2, \tag{3.3}$$

where m is the trace of M and denote by

$$\|h\|^2 = \sum_{i,j=1}^n g(h(e_i, e_j), h(e_i, e_j)).$$

From (3.3), we get

$$\begin{aligned} \frac{1}{4}n^2 \|H\|^2 &= \tau(p) - \frac{n(n-1)c}{2} + (n-1)^2 cm \\ &+ \frac{1}{4} \sum_{r=n+1}^m (h_{11}^r - h_{22}^r - \dots - h_{nn}^r)^2 + \sum_{r=n+1}^m \sum_{j=2}^n (h_{1j}^r)^2 \\ &- \sum_{r=n+1}^m \sum_{2 \leq i < j \leq n} (h_{ii}^r h_{jj}^r - (h_{ij}^r)^2), \end{aligned} \tag{3.4}$$

where

$$h_{ij}^r = g(h(e_i, e_j), e_r).$$

Using (2.7) and (2.15) we also have

$$\begin{aligned} \sum_{r=n+1}^m \sum_{2 \leq i < j \leq n} (h_{ii}^r h_{jj}^r - (h_{ij}^r)^2) &= \sum_{2 \leq i < j \leq n} K_{ij} - \sum_{2 \leq i < j \leq n} \tilde{K}_{ij} \\ &= \sum_{2 \leq i < j \leq n} K_{ij} - (n-1)(n-2)c \left(\frac{1}{2} - m + M(e_1, e_1) \right). \end{aligned} \tag{3.5}$$

From (3.4) and (3.5), we obtain

$$\begin{aligned} Ric(e_1) &= \frac{1}{4}n^2 \|H\|^2 - (n-1)c(1-m+(n-2)M(e_1, e_1)) \\ &- \sum_{r=n+1}^m \sum_{j=2}^n (h_{1j}^r)^2 - \frac{1}{4} \sum_{r=n+1}^m (h_{11}^r - h_{22}^r - \dots - h_{nn}^r)^2. \end{aligned} \tag{3.6}$$

If we choose $e_1 = X$ as any unit vector of $T_p M^n$ in the above equation, one obtains (3.1).

Taking into consideration equation (3.6) and $X = e_1$, the equality case of (3.1) holds if and only if

$$h_{12}^r = h_{13}^r = \dots = h_{1n}^r = 0 \text{ and } h_{11}^r = h_{22}^r + \dots + h_{nn}^r, \quad r \in \{n+1, \dots, m\} \tag{3.7}$$

which shows that (3.2) holds.

We now suppose that the equality case of (3.1) holds for all unit vector $X \in T_p M^n$. Then, in view of (3.7), for each $r \in \{n+1, \dots, m\}$ we have $i \in \{1, \dots, n\}$,

$$h_{ij}^r = 0, \quad i \neq j \tag{3.8}$$

$$2h_{ii}^r = h_{11}^r + h_{22}^r + \dots + h_{nn}^r, \quad i \in \{1, \dots, n\}. \tag{3.9}$$

From (3.9), we have $2h_{11}^r = 2h_{22}^r = \dots = 2h_{nn}^r = h_{11}^r + h_{22}^r + \dots + h_{nn}^r$, which implies that

$$(n-2)(h_{11}^r + h_{22}^r + \dots + h_{nn}^r) = 0. \tag{3.10}$$

Thus, either $h_{11}^r + h_{22}^r + \dots + h_{nn}^r = 0$ or $n = 2$. If $h_{11}^r + h_{22}^r + \dots + h_{nn}^r = 0$, then in view of (3.9), we get $h_{ii}^r = 0$ for all $i \in \{1, \dots, n\}$. This together with (3.8) gives $h_{ij}^r = 0$ for all $i, j \in \{1, \dots, n\}$ and $r \in \{n+1, \dots, m\}$, that is, p is a totally geodesic point. If $n = 2$, then from (3.9) $2h_{11}^r = 2h_{22}^r = h_{11}^r + h_{22}^r$, which shows that p is a totally umbilical point. The proof of the converse part is straightforward. \square

Corollary 3.1. *If $H(p) = 0$, then a unit tangent vector X at p satisfies the equality case of (3.1) if and only if $X \in N(p)$.*

Theorem 3.2. *Let M^n be an n -dimensional submanifold of an m -dimensional real space form $\tilde{M}(c)$ of constant sectional curvature c endowed with Ricci quarter-symmetric metric connection $\tilde{\nabla}$*

$$\tau(p) \leq \frac{(n-1)}{2} \left(n \|H\|^2 + nc - 2c(n-1)m \right). \tag{3.11}$$

Equality case of (3.11) holds at $p \in M^n$ if and only if p is a totally umbilical point.

Proof. Let $p \in M^n$ and $\{e_1, \dots, e_n\}$ be orthonormal basis of $T_p M^n$. The relation (3.3) is equivalent to

$$n^2 \|H\|^2 = 2\tau(p) + \|h\|^2 + (n-1)c(2(n-1)m-n). \tag{3.12}$$

We choose an orthonormal basis $\{e_1, \dots, e_n, e_{n+1}, \dots, e_m\}$ at p such that e_{n+1} is parallel to the mean curvature vector $H(p)$ and e_1, \dots, e_n diagonalize the shape operator $A_{e_{n+1}}$. Then the shape operators take the forms

$$A_{e_{n+1}} = \begin{bmatrix} a_1 & 0 & \cdot & \cdot & \cdot & 0 \\ 0 & a_2 & \cdot & \cdot & \cdot & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ 0 & 0 & \cdot & \cdot & \cdot & a_n \end{bmatrix} \tag{3.13}$$

$$A_{e_r} = (h_{ij}^r), \quad i, j = 1, \dots, n; \quad r = n+2, \dots, m, \quad \text{trace} A_{e_r} = 0. \tag{3.14}$$

From (3.12), we get

$$n^2 \|H\|^2 = 2\tau(p) + \sum_{i=1}^n a_i^2 + \sum_{r=n+2}^m \sum_{i,j=1}^n (h_{ij}^r)^2 + (n-1)c(2(n-1)m-n). \tag{3.15}$$

On the other hand, since

$$0 \leq \sum_{i<j} (a_i - a_j)^2 = (n-1) \sum_i a_i^2 - 2 \sum_{i<j} a_i a_j \tag{3.16}$$

we obtain

$$n^2 \|H\|^2 = \left(\sum_{i=1}^n a_i \right)^2 = \sum_{i=1}^n a_i^2 + 2 \sum_{i<j} a_i a_j \leq n \sum_{i=1}^n a_i^2 \tag{3.17}$$

which implies

$$\sum_{i=1}^n a_i^2 \geq n \|H\|^2. \tag{3.18}$$

So from (3.15) and (3.18), we have

$$n^2 \|H\|^2 \geq 2\tau(p) + n \|H\|^2 + \sum_{r=n+2}^m \sum_{i,j=1}^n (h_{ij}^r)^2 + (n-1)c(2(n-1)m-n). \tag{3.19}$$

If the equality case of (3.11) holds, then from (3.16) and (3.19) it follows that

$$a_1 = a_2 = \dots = a_n \quad \text{and} \quad A_{e_r} = 0, \quad r = n+2, \dots, m. \tag{3.20}$$

Therefore, p is a totally umbilical point. The converse is straightforward. □

Theorem 3.3. *Let M^n be an n -dimensional submanifold of an m -dimensional real space form $\widetilde{M}(c)$ of constant sectional curvature c endowed with Ricci quarter-symmetric metric connection $\widetilde{\nabla}$. Then we have*

$$\theta_k(p) \leq \|H\|^2 + c \left(2 - \frac{4(n-1)m}{n} \right). \tag{3.21}$$

Proof. Let $\{e_1, \dots, e_n\}$ be an orthonormal basis of $T_p M^n$. Denote by $L_{i_1 \dots i_k}$ the k -plane section spanned by $\{e_{i_1}, \dots, e_{i_k}\}$. Using the definitions of the ricci and scalar curvatures, we have

$$\tau(L_{i_1 \dots i_k}) = \frac{1}{2} \sum_{i \in \{i_1, \dots, i_k\}} Ric_{L_{i_1 \dots i_k}}(e_i), \tag{3.22}$$

$$\tau(p) = \frac{1}{C_{n-2}^{k-2}} \sum_{1 \leq i_1 < \dots < i_k \leq n} \tau(L_{i_1 \dots i_k}). \tag{3.23}$$

From (2.10), (3.22) and (3.23), we get

$$\tau(p) \geq \frac{n(n-1)}{2} \theta_k(p). \tag{3.24}$$

Using (3.11) and (3.24) we obtain (3.21). □

Lemma 3.1. If $n > k \geq 2$ and a_1, \dots, a_n, a are real numbers such that

$$\left(\sum_{i=1}^n a_i\right)^2 = (n - k + 1) \left(\sum_{i=1}^n a_i^2 + a\right) \tag{3.25}$$

then

$$2 \sum_{1 \leq i < j \leq k} a_i a_j \geq a \tag{3.26}$$

with equality holding if and only if

$$a_1 + a_2 + \dots + a_k = a_{k+1} = \dots = a_n. \tag{3.27}$$

Theorem 3.4. Let M^n be an n -dimensional submanifold of an m -dimensional real space form $\widetilde{M}(c)$ of constant sectional curvature c endowed with Ricci quarter-symmetric metric connection $\widetilde{\nabla}$. Then, for each point $p \in M^n$ and each k -plane section $\Pi_k \subset TpM^n$ ($n > k \geq 2$), we have

$$\begin{aligned} \tau(p) - \tau(\pi_k) \leq & (n - k) \left[\frac{n^2}{2(n - k + 1)} \|H\|^2 - \frac{(n + k - 1)}{2} c - (n - 1)cm \right] \\ & - (n - 1)(k - 1)c \operatorname{trace}(m_{\pi_k^\perp}). \end{aligned} \tag{3.28}$$

The equality case of (3.28) holds at $p \in M^n$ if and only if there exist an orthonormal basis $\{e_1, \dots, e_n\}$ of TpM^n and an orthonormal basis $\{e_{n+1}, \dots, e_m\}$ of $T_p^\perp M^n$ such that (a) $\Pi_k = \operatorname{Span}\{e_1, \dots, e_k\}$ and (b) the forms of shape operators A_{e_r} , $r = n + 1, \dots, m$, take the forms

$$\begin{aligned} A_{e_{n+1}} &= \begin{bmatrix} h_{11}^{n+1} & 0 & \cdot & \cdot & \cdot & 0 \\ 0 & h_{22}^{n+1} & \cdot & \cdot & \cdot & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot & 0 \\ 0 & 0 & \cdot & \cdot & \cdot & h_{kk}^{n+1} \\ & & & & 0 & \left(\sum_{i=1}^k h_{ii}^{n+1}\right) I_{n-k} \end{bmatrix}, \tag{3.29} \\ A_{e_r} &= \begin{bmatrix} h_{11}^r & h_{12}^r & \cdot & \cdot & \cdot & h_{1k}^r \\ h_{12}^r & h_{22}^r & \cdot & \cdot & \cdot & h_{2k}^r \\ \cdot & \cdot & \cdot & \cdot & \cdot & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot & 0 \\ h_{1k}^r & h_{2k}^r & \cdot & \cdot & \cdot & -\sum_{i=1}^{k-1} h_{ii}^r \\ & & & & 0 & 0_{n-k} \end{bmatrix}, \quad r \in \{n + 2, \dots, m\}. \end{aligned} \tag{3.30}$$

Proof. Let $\Pi_k \subset TpM^n$ be a k -plane section. We choose an orthonormal basis $\{e_1, \dots, e_n\}$ for TpM^n and $\{e_{n+1}, \dots, e_m\}$ for the normal space $T_p^\perp M^n$ at p such that $\Pi_k = \operatorname{Span}\{e_1, \dots, e_k\}$, the mean curvature vector H is in the direction of the normal vector to e_{n+1} and e_1, \dots, e_n diagonalize the shape operator $A_{e_{n+1}}$. Then the shape operators take the forms (3.13) and (3.14). We rewrite (3.3) as

$$\left(\sum_{i=1}^n h_{ii}^{n+1}\right)^2 = (n - k + 1) \left(\sum_{i=1}^n (h_{ii}^{n+1})^2 + \sum_{r=n+2}^m \sum_{i,j=1}^n (h_{ij}^r)^2 + \epsilon\right), \tag{3.31}$$

where

$$\epsilon = 2\tau(p) - n(n - 1)c + 2(n - 1)^2 cm - \frac{n^2(n - k)}{(n - k + 1)} \|H\|^2. \tag{3.32}$$

Applying Lemma 3.1 in (3.31), we get

$$2 \sum_{1 \leq i < j \leq k} h_{ii}^{n+1} h_{jj}^{n+1} \geq \epsilon + \sum_{r=n+2}^m \sum_{i,j=1}^n (h_{ij}^r)^2. \tag{3.33}$$

From equation (2.7) and (2.15) it also follows that

$$\begin{aligned} \tau(\pi_k) &= \frac{k(k-1)c}{2} - (n-1)(k-1)c \sum_{i=1}^k M(e_i, e_i) + \sum_{1 \leq i < j \leq k} h_{ii}^{n+1} h_{jj}^{n+1} \\ &+ \sum_{r=n+2}^m \sum_{1 \leq i < j \leq k} (h_{ii}^r h_{jj}^r - (h_{ij}^r)^2). \end{aligned} \tag{3.34}$$

Using (3.33) and (3.34) we get

$$\begin{aligned} \tau(\pi_k) &\geq \frac{k(k-1)c}{2} - (n-1)(k-1)c \sum_{i=1}^k M(e_i, e_i) + \frac{1}{2}\epsilon \\ &+ \frac{1}{2} \sum_{r=n+2}^m (h_{11}^r + h_{22}^r + \dots + h_{kk}^r)^2 + \frac{1}{2} \sum_{r=n+2}^m \sum_{i,j>k}^n (h_{ij}^r)^2 \\ &+ \sum_{r=n+2}^m \sum_{j>k}^n ((h_{1j}^r)^2 + (h_{2j}^r)^2 + \dots + (h_{kj}^r)^2) \end{aligned} \tag{3.35}$$

or

$$\tau(\pi_k) \geq \frac{k(k-1)c}{2} - (n-1)(k-1)c \sum_{i=1}^k M(e_i, e_i) + \frac{1}{2}\epsilon. \tag{3.36}$$

We remark that

$$M(e_1, e_1) + M(e_2, e_2) + \dots + M(e_k, e_k) = m - \text{trace}(m_{|\pi_k^\perp}). \tag{3.37}$$

From (3.32), (3.36) and (3.37) we obtain

$$\begin{aligned} \tau(\pi_k) &\geq (n-k) \left(\frac{(n+k-1)}{2}c + (n-1)cm \right) + \tau(p) \\ &- \frac{n^2(n-k)}{2(n-k+1)} \|H\|^2 + (n-1)(k-1)c \text{trace}(m_{|\pi_k^\perp}) \end{aligned} \tag{3.38}$$

which proves the inequality case of (3.28). □

If the equality case of (3.28) holds, then the inequalities given by (3.33) and (3.36) become equalities. In this case, for $r = n + 2, \dots, m$ we have

$$h_{1j}^{n+1} = h_{2j}^{n+1} = \dots = h_{kj}^{n+1} = 0, \quad j = k + 1, \dots, n, \tag{3.39}$$

$$h_{ij}^r = 0, \quad i, j = k + 1, \dots, n, \tag{3.40}$$

$$h_{11}^r + h_{22}^r + \dots + h_{kk}^r = 0. \tag{3.41}$$

Applying Lemma 3.1 we also have

$$h_{11}^{n+1} + h_{22}^{n+1} + \dots + h_{kk}^{n+1} = h_{ll}^{n+1}, \quad l = k + 1, \dots, n. \tag{3.42}$$

Thus, after choosing a suitable orthonormal basis $\{e_1, \dots, e_m\}$, the shape operator of M^n takes the form given by (3.29) and (3.30). The converse is easy to follow.

By Theorem 3.4 we get the following corollary.

Corollary 3.2. *Let M^n , $n \geq 3$, be an n -dimensional submanifold of an m -dimensional real space form $\widetilde{M}(c)$ of constant sectional curvature c endowed with Ricci quarter-symmetric metric connection $\widetilde{\nabla}$. Then, for each point $p \in M^n$ and each 2-plane section $\Pi_2 \subset T_p M^n$, we have*

$$\begin{aligned} \delta_M &\leq (n-2) \left[\frac{n^2}{2(n-1)} \|H\|^2 - \frac{(n+1)}{2}c - (n-1)cm \right] \\ &- (n-1)c \text{trace}(m_{|\pi_2^\perp}). \end{aligned} \tag{3.43}$$

The equality case of (3.43) holds at $p \in M^n$ if and only if there exist an orthonormal basis $\{e_1, \dots, e_2\}$ of $T_p M^n$ and an orthonormal basis $\{e_{n+1}, \dots, e_m\}$ of $T_p^\perp M^n$ such that (a) $\Pi_2 = \text{Span}\{e_1, e_2\}$ and (b) the forms of shape operators A_{e_r} , $r = n + 1, \dots, m$, become

$$A_{e_{n+1}} = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & (a+b)I_{n-2} \end{bmatrix}, A_{e_r} = \begin{bmatrix} c_r & d_r & 0 \\ d_r & -c_r & 0 \\ 0 & 0 & 0_{n-2} \end{bmatrix}, r = n + 2, \dots, m. \quad (3.44)$$

References

- [1] Chen, B. Y., Mean curvature and shape operator of isometric immersion in real space forms. *Glasgow Mathematic Journal* 38 (1996), 87-97.
- [2] Chen, B. Y., Relation between Ricci curvature and shape operator for submanifolds with arbitrary codimension. *Glasgow Mathematic Journal* 41 (1999), 33-41.
- [3] Chen, B. Y., Some pinching and classification theorems for minimal submanifolds. *Arch. math* (Basel) 60 (1993), no. 6, 568-578.
- [4] Chen, B. Y., A Riemannian invariant for submanifolds in space forms and its applications. *Geometry and Topology of submanifolds VI*. (Leuven, 1993/Brussels, 1993). (NJ:Word Scientific Publishing, River Edge). 1994, pp. 58-81, no. 6, 568-578.
- [5] Chen, B. Y., A general optimal inequality for arbitrary Riemannian submanifolds. *J. Ineq. Pure Appl. Math* 6 (2005), no. 3, Article 77, 1-11.
- [6] Gülbahar, M., Kılıç, E., Keleş, S. and Tripathi, M. M., Some basic inequalities for submanifolds of nearly quasi-constant curvature manifolds. *Differential Geometry-Dynamical Systems*. 16 (2014), 156-167.
- [7] Hong, S. and Tripathi, M. M., On Ricci curvature of submanifolds. *Int J. Pure Appl. Math. Sci.* 2 (2005), no.2, 227-245.
- [8] Kamilya, D and De, U. C., Some properties of a Ricci quarter-symmetric metric connection in a Riemannian manifold. *Indian J. Pure and Appl. Math* 26 (1995), no. 1, 29-34.
- [9] Liu, X. and Zhou, J., On Ricci curvature of certain submanifolds in cosymplectic form. *Sarajevo J. Math* 2 (2006), no.1, 95-106.
- [10] Mihai, A. and Özgür, C., Chen inequalities for submanifolds of real space form with a semi-symmetric metric connection. *Taiwanese Journal of Mathematics* 14 (2010), no. 4, 1465-1477.
- [11] Mishra, R. S. and Pandey, S. N., On quarter symmetric metric F-connections. *Tensor (N.S.)* 34 (1980), no. 1, 1-7.
- [12] Rastogi, S. C., On quarter-symmetric metric connection. *C. R. Acad. Bulgare Sci* 31 (1978), no. 7, 811-814.
- [13] Tripathi, M. M., Improved Chen-Ricci inequality for curvature-like tensor and its applications. *Differential Geom. Appl.* 29 (2011), 685-698.

Affiliations

NERGİZ (ÖNEN) POYRAZ

ADDRESS: Cukurova University, Department of Mathematics,
01330, Adana-Turkey.

E-MAIL: nonen@cu.edu.tr

ORCID ID: orcid.org/0000-0002-8110-712X

HALİL İBRAHİM YOLDAŞ

ADDRESS: Mersin University, Department of Mathematics,
33343, Mersin-Turkey.

E-MAIL: hibrahimyoldas@mersin.edu.tr

ORCID ID: orcid.org/0000-0002-3238-6484